

**Section 3.5**

**Graphing Techniques;  
Transformations**

# OBJECTIVE 1



**Graph Functions Using Vertical and Horizontal Shifts**


# Exploration

On the same screen, graph each of the following functions:

$$Y_1 = x^2$$

$$Y_2 = x^2 + 2$$

$$Y_3 = x^2 - 2$$



If a positive real number  $k$  is added to the outputs of a function  $y = f(x)$ , the graph of the new function  $y = f(x) + k$  is the graph of  $f$  **shifted vertically up**  $k$  units.

If a positive real number  $k$  is subtracted from the outputs of a function  $y = f(x)$ , the graph of the new function  $y = f(x) - k$  is the graph of  $f$  **shifted vertically down**  $k$  units.

**EXAMPLE****Vertical Shift**


Use the graph of  $f(x) = x^2$  to obtain the graph of the following:

(a)  $g(x) = x^2 + 2$

(b)  $h(x) = x^2 - 2$

# Exploration

On the same screen, graph each of the following functions:



$$Y_1 = x^2$$

$$Y_2 = (x - 3)^2$$

$$Y_3 = (x + 2)^2$$

If the argument  $x$  of a function  $f$  is replaced by  $x - h$ ,  $h > 0$ , the graph of the new function  $y = f(x - h)$  is the graph of  $f$  **shifted horizontally right**  $h$  units. If the argument  $x$  of a function  $f$  is replaced by  $x + h$ ,  $h > 0$ , the graph of the new function  $y = f(x + h)$  is the graph of  $f$  **shifted horizontally left**  $h$  units..

## EXAMPLE

### Combining Vertical and Horizontal Shifts

Graph the function  $f(x) = (x - 2)^2 - 3$

# OBJECTIVE 2

 **2 Graph Functions Using Compressions and Stretches**


## Exploration

On the same screen, graph each of the following functions:

$$Y_1 = |x|$$

$$Y_2 = 2|x|$$

$$Y_3 = \frac{1}{2}|x|$$



$$Y_1 = |x|$$

$$Y_2 = 2|x|$$

$$Y_3 = \frac{1}{2}|x|$$

X	Y1	Y2
-2	2	4
-1	1	2
0	0	0
1	1	2
2	2	4
3	3	6
4	4	8

$Y_2 = 2\text{abs}(X)$

X	Y1	Y3
-2	2	1
-1	1	.5
0	0	0
1	1	.5
2	2	1
3	3	1.5
4	4	2

$Y_3 = .5\text{abs}(X)$



## Exploration


On the same screen, graph each of the following functions:

$$Y_1 = |x|$$
$$Y_2 = 2|x|$$
$$Y_3 = \frac{1}{2}|x|$$

$$Y_1 = |x|$$

$$Y_2 = 2|x|$$

$$Y_3 = \frac{1}{2}|x|$$




When the right side of a function  $y = f(x)$  is multiplied by a positive number  $a$ , the graph of the new function  $y = af(x)$  is obtained by multiplying each  $y$ -coordinate on the graph of  $y = f(x)$  by  $a$ . The new graph is a **vertically compressed** (if  $0 < a < 1$ ) or a **vertically stretched** (if  $a > 1$ ) version of the graph of  $y = f(x)$ .

# Exploration

On the same screen, graph each of the following functions:

$$Y_1 = f(x) = \sqrt{x} \quad Y_2 = f(2x) = \sqrt{2x} \quad Y_3 = f\left(\frac{1}{2}x\right) = \sqrt{\frac{1}{2}x} = \sqrt{\frac{x}{2}}$$



X	Y <sub>1</sub>	Y <sub>2</sub>
0	0	0
.5	.70711	1
1	1	1.4142
2	1.4142	2
4	2	2.8284
4.5	2.1213	3
9	3	4.2426

$Y_2 = \sqrt{2X}$


X	Y <sub>1</sub>	Y <sub>3</sub>
0	0	0
1	1	.70711
2	1.4142	1
4	2	1.4142
8	2.8284	2
9	3	2.1213
18	4.2426	3

$Y_3 = \sqrt{X/2}$

## Exploration

On the same screen, graph each of the following functions:

$$Y_1 = f(x) = \sqrt{x} \quad Y_2 = f(2x) = \sqrt{2x} \quad Y_3 = f\left(\frac{1}{2}x\right) = \sqrt{\frac{1}{2}x} = \sqrt{\frac{x}{2}}$$






If the argument  $x$  of a function  $y = f(x)$  is multiplied by a positive number  $a$ , the graph of the new function  $y = f(ax)$  is obtained by multiplying each  $x$ -coordinate of  $y = f(x)$  by  $\frac{1}{a}$ . A **horizontal compression** results if  $a > 1$ , and a **horizontal stretch** occurs if  $0 < a < 1$ .

## EXAMPLE

### Graphing Using Stretches and Compressions

The graph of  $y = f(x)$  is given.  
Use this graph to find the graphs of

(a)  $y = 2f(x)$       (b)  $y = f(3x)$



# OBJECTIVE 3




**Graph Functions Using Reflections about  
the  $x$ -Axis or  $y$ -Axis**

# Exploration

Reflection about the x-axis:

(a) Graph  $Y_1 = x^2$ , followed by  $Y_2 = -x^2$ .

X	Y <sub>1</sub>	Y <sub>2</sub>
-3	9	-9
-2	4	-4
-1	1	-1
0	0	0
1	1	-1
2	4	-4
3	9	-9
Y <sub>2</sub> = -X <sup>2</sup>		






# Exploration

(b) Graph  $Y_1 = |x|$ , followed by  $Y_2 = -|x|$ .

X	Y <sub>1</sub>	Y <sub>2</sub>
-3	3	-3
-2	2	-2
-1	1	-1
0	0	0
1	1	-1
2	2	-2
3	3	-3

$Y_2 = -\text{abs}(X)$




# Exploration

(c) Graph  $Y_1 = x^2 - 4$ , followed by  $Y_2 = -(x^2 - 4) = -x^2 + 4$ .

X	Y <sub>1</sub>	Y <sub>2</sub>
-3	5	-5
-2	0	0
-1	-3	3
0	-4	4
1	-3	3
2	0	0
3	5	-5

Y<sub>2</sub> = -X<sup>2</sup> + 4



When the right side of the function  $y = f(x)$  is multiplied by  $-1$ , the graph of the new function  $y = -f(x)$  is the **reflection about the x-axis** of the graph of the function  $y = f(x)$ .




# Exploration

Reflection about the  $y$ -axis:

(a) Graph  $Y_1 = \sqrt{x}$ , followed by  $Y_2 = \sqrt{-x}$ .

X	Y <sub>1</sub>	Y <sub>2</sub>
-3	ERROR	1.7321
-2	ERROR	1.4142
-1	ERROR	1
0	0	0
1	1	ERROR
2	1.4142	ERROR
3	1.7321	ERROR

$Y_2 = \sqrt{(-X)}$




# Exploration

(b) Graph  $Y_1 = x + 1$ , followed by  $Y_2 = -x + 1$ .

X	Y <sub>1</sub>	Y <sub>2</sub>
-3	-2	4
-2	-1	3
-1	0	2
0	1	1
1	2	0
2	3	-1
3	4	-2

Y<sub>2</sub> = -X + 1




# Exploration

(c) Graph  $Y_1 = x^4 + x$ , followed by  $Y_2 = (-x)^4 + (-x) = x^4 - x$ .

X	Y <sub>1</sub>	Y <sub>2</sub>
-3	78	84
-2	14	18
-1	0	2
0	0	0
1	2	0
2	18	14
3	84	78

Y<sub>2</sub> = X<sup>4</sup> - X



When the graph of the function  $y = f(x)$  is known, the graph of the new function  $y = f(-x)$  is the **reflection about the y-axis** of the graph of the function  $y = f(x)$ .



## Summary of Graphing Techniques

### To Graph:

### Draw the Graph of $f$ and:

### Functional Change to $f(x)$

#### Vertical shifts

$$y = f(x) + k, \quad k > 0$$

Raise the graph of  $f$  by  $k$  units.

Add  $k$  to  $f(x)$ .

$$y = f(x) - k, \quad k > 0$$

Lower the graph of  $f$  by  $k$  units

Subtract  $k$  from  $f(x)$ .

#### Horizontal shifts

$$y = f(x + h), \quad h > 0$$

Shift the graph of  $f$  to the left  $h$  units.

Replace  $x$  by  $x + h$ .

$$y = f(x - h), \quad h > 0$$

Shift the graph of  $f$  to the right  $h$  units.

Replace  $x$  by  $x - h$ .



# Summary of Graphing Techniques

**To Graph:**

**Draw the Graph of  $f$  and:**

**Functional Change to  $f(x)$**

## Compressing or stretching

$$y = af(x), \quad a > 0$$

Multiply each  $y$ -coordinate of  $y = f(x)$  by  $a$ .

Stretch the graph of  $f$  vertically if  $a > 1$ .

Compress the graph of  $f$  vertically if  $0 < a < 1$ .

Multiply  $f(x)$  by  $a$ .

$$y = f(ax), \quad a > 0$$

Multiply each  $x$ -coordinate of  $y = f(x)$  by  $\frac{1}{a}$ .

Stretch the graph of  $f$  horizontally if  $0 < a < 1$ .

Compress the graph of  $f$  horizontally if  $a > 1$ .

Replace  $x$  by  $ax$ .



## Summary of Graphing Techniques

**To Graph:**

**Draw the Graph of  $f$  and:**

**Functional Change to  $f(x)$**

**Reflection about the  $x$ -axis**

$$y = -f(x)$$

**Reflection about the  $y$ -axis**

$$y = f(-x)$$

Reflect the graph of  $f$  about the  $x$ -axis.

Reflect the graph of  $f$  about the  $y$ -axis.

Multiply  $f(x)$  by  $-1$ .

Replace  $x$  by  $-x$ .

## EXAMPLE

### Determining the Function Obtained from a Series of Transformations

Find the function that is finally graphed after the following three transformations are applied to the graph of  $y = \sqrt{x}$

1. Shift right 1 unit.
2. Shift down 3 units.
3. Reflect about the  $x$ -axis.

**EXAMPLE****Combining Graphing Procedures**

Graph the function  $f(x) = -2|x + 1| - 3$ .

Find the domain and the range of  $f$ .



**EXAMPLE****Combining Graphing Procedures**

Graph the function  $f(x) = \frac{2}{x+3} + 4$ .

Find the domain and the range of  $f$ .