

## **Section 8.3**

# **Trigonometric Identities**

Two functions  $f$  and  $g$  are said to be **identically equal** if

$$f(x) = g(x)$$

for every value of  $x$  for which both functions are defined. Such an equation is referred to as an **identity**. An equation that is not an identity is called a **conditional equation**.

## Quotient Identities

$$\tan \theta = \frac{\sin \theta}{\cos \theta} \quad \cot \theta = \frac{\cos \theta}{\sin \theta}$$

## Reciprocal Identities

$$\csc \theta = \frac{1}{\sin \theta} \quad \sec \theta = \frac{1}{\cos \theta} \quad \cot \theta = \frac{1}{\tan \theta}$$

## Pythagorean Identities

$$\sin^2 \theta + \cos^2 \theta = 1 \quad \tan^2 \theta + 1 = \sec^2 \theta$$
$$\cot^2 \theta + 1 = \csc^2 \theta$$

## Even-Odd Identities

$$\sin(-\theta) = -\sin \theta \quad \cos(-\theta) = \cos \theta \quad \tan(-\theta) = -\tan \theta$$
$$\csc(-\theta) = -\csc \theta \quad \sec(-\theta) = \sec \theta \quad \cot(-\theta) = -\cot \theta$$

# OBJECTIVE 1

- ✓ Use Algebra to Simplify Trigonometric Expressions

## EXAMPLE

### Using Algebraic Techniques to Simplify Trigonometric Expressions

(a) Simplify  $\frac{\tan \theta}{\sec \theta}$  by rewriting each trigonometric function in terms of sine and cosine functions.

(b) Show that  $\frac{\sin \theta}{1 + \cos \theta} = \frac{1 - \cos \theta}{\sin \theta}$  by multiplying the numerator and denominator by  $1 - \cos \theta$

(c) Simplify  $\frac{1}{1 - \sin u} + \frac{1}{1 + \sin u}$  by rewriting the expression over a common denominator.

(d) Simplify  $\frac{1 - \cos^2 v}{\sin v + \cos v \sin v}$  by factoring.

# OBJECTIVE 2

 **2 Establish Identities**

**EXAMPLE****Establishing an Identity**

Establish the identity:  $\sin \theta (\cot \theta + \tan \theta) = \sec \theta$



**EXAMPLE****Establishing an Identity**

Establish the identity:  $\csc \theta - \cot \theta = \frac{\sin \theta}{1 + \cos \theta}$

**EXAMPLE****Establishing an Identity**

Establish the identity:  $\frac{\sin^2 \theta - \tan \theta}{\cos^2 \theta - \cot \theta} = \tan^2 \theta$

**EXAMPLE****Establishing an Identity**

Establish the identity:  $\frac{\cos \theta}{1 - \sin \theta} = \frac{1 + \sin \theta}{\cos \theta}$

**EXAMPLE****Establishing an Identity**

Establish the identity:  $\cot^2 \theta = \frac{\csc \theta - \sin \theta}{\sin \theta}$

**EXAMPLE****Establishing an Identity**

Establish the identity:  $1 - \csc \theta \sin^3 \theta = \cos^2 \theta$

## **Guidelines for Establishing Identities**

- 1.** It is almost always preferable to start with the side containing the more complicated expression.
- 2.** Rewrite sums or differences of quotients as a single quotient.
- 3.** Sometimes rewriting one side in terms of sines and cosines only will help.
- 4.** Always keep your goal in mind. As you manipulate one side of the expression, you must keep in mind the form of the expression on the other side.